

Diversity and Nestedness pattern of adult Odonata assemblages around Hadhinaru lake of Mysore, Karnataka

Dr. Shankar Hosmani

Professor and Head, Department of Biotechnology, SBRR Mahajana First Grade College,
Jayalakshmipuram, Mysore
Email: prof_sph@yahoo.co.in

ABSTRACT

Diversity and nestedness assemblages of adult insect Odonata around Hadhinaru Lake of Mysore was done for a period of three years. 23 species belonging to Dragonflies (Anisoptera) and Damselflies (Zygoptera) were recorded. Their diversity was not quite significant. The order of nestedness of species showed a hot matrix (36.7⁰), with a maximum fill of 46.3%. Nested species were low while idiosyncratic species were abundant. Species appearance and disappearance was fairly common which may be attributed to the human disturbances in the lake. Odonates can help control small insects like mosquitoes and hence their conservation is of importance. The nestedness software serves as a handy tool in monitoring insect biodiversity.

Keywords: Dragonflies, Damselflies, Nestedness, Idiosyncratic, Temperature

INTRODUCTION

Odonata is an order of insects, encompassing Dragonflies (Anisoptera) and Damselflies (Zygoptera). About 5900 species have been described in this order. Most of these are regarded as beneficial because they feed on small insects like mosquitoes, but act as pests to beekeepers, since they eat honeybees also. Secondly they transmit *Prosthogoniums pellucidus*, a parasitic flatworm of poultry and hence are a matter of concern. Their scientific classification includes Kingdom: Animalia; Phylum: Arthropoda; Class: Insecta; Subclass: Pterygota; Infra class Paleoptera; Sub Order Odonatoptera and Order Odonata (*Fabricus, 1973*). The suborders include Epirocta (dragonflies) and Zygoptera (damselflies). These insects characteristically have large rounded heads covered mostly by well developed compound eyes, legs that facilitate catching prey, two pairs of long transparent wings that move independently and have an elongated abdomen. Dragonflies differ from damselflies in many recognizable traits. Dragonfly eyes occupy much of the animals head touching each other across the face, while in damselflies there is typically a gap between the eyes. Odonates are aquatic as juveniles and are often described as aquatic insects.

Nestedness is a measure of order in ecological systems referring to the order in which the number of species is related to the area or other factors. A challenge before ecologists is to

understand the structure and dynamics of biological communities and their relation to environmental variables. Werner and Peacor (1990) and Levins(1992) are of the opinion that understanding the community and its structure is a fundamental aspect of conservation biology. Spatial patterns for insect diversity have not been dealt with in detail and studies on Odonata diversity are very meagre. Soinen *et al.* (2009) says that dispersion and migration of communities at regional levels need to be stressed. Studies on nestedness assemblages, other than insects are those of Wright *et al.* (1998), Brown (1995) and Cook and Quinn (1995). However species assemblage patterns of tropical fresh water lakes in India are absolutely lacking. An attempt to study the nestedness pattern of Odonata insects around a very large lake (Hadhinaru Lake) of Mysore has been done. Nestedness assemblages are important in lake conservation strategies.

Material and Methods

Hadhinaru Lake is located at 12⁰2' north latitude and 76⁰4' east longitude at an altitude of 753 meters above MSL and is situated 16 kms away from Mysore city (Karnataka). It has an independent catchment area of 9.57sq.km with water spread over 10.10 hectares having a live capacity of 54.43 mcft. The maximum depth of the lake when full is five meters. It is important from the Social-Economic-Natural-Cultural-Lake-Ecosystem (SENACLE) of the area.

Adult Odonata sampling was done by direct counts made while observing habitats on hourly basis in the early, mid and late hours of the day during suitable flight conditions(during low wind, warm and sunny weather). An aerial net (30x15x60 cms) was used to catch the insects. Disturbance by a surveyor caused a flight response making sedentary insects visible (Moore, 1953). Counts were standardized based on time and catch per unit effort (CPUE). The insects were let free later (Foote *et al.*2005), Conard *et al.* 1999).Odonata naiads were however not accounted in the present study. The insects were identified using taxonomic keys of Fraser (1933-1934), Ward and Whipple (1959), Melamby(1963),Pennak(1978), Tonapi(1980), Emiliyamma *et al.*(2005) and Subramanian(2005).


RESULTS AND DISCUSSION

In the present investigation, 23 species, both males and females were recorded separately wherever identifiable. Ten species of the order Anisoptera, Family Libellulidae, one species each of Ashnidae and Gomphidae and four species of the order Zygoptera, Family Coenagrionidae and one species of Lestidae were found to be of common occurrence around the lake. The scientific and common names of the insects are presented in Table 1.

Nestedness analysis was carried out using the presence-absence matrix (Matrix having sites in rows and species in columns with "1" for presence and "0" for absence was considered for the insect diversity. The degree of nestedness was calculated using the "Nestedness Calculator" (Atmar and Patterson, 1993, 1995). The metric "T" of the Nestedness Calculator measures the extent of unexpected presences and absences in a maximally packed matrix, where sites are arranged in rows in descending order according to their species richness and the species are arranged in columns according to the number of sites in which they occur. The metric "T"⁰ for a set of perfectly nested assemblages and "T"¹⁰⁰ for a completely disordered matrix is expressed. The null hypothesis that "T" is not lower (more nested) than expected by chance was tested using Monte Carlo permutations of 100 counts (Heino and

Moutka, 2005). The reorganized species vector is presented in Table 2. Different species recorded are shown in Plate 1 & 2 and the nestedness matrix is presented in Fig. 1.

Table 1. Species index of Odonata

Order/ Suborder	Family	Scientific name	Common name	
Odonata: Anisoptera	Libellulidae	<i>Acisoma panorpoides</i> Ramb	Trumpet Tail	
 <p>Damselflies</p> <p>Dragonflies</p>		<i>Brachydipax sorbina</i> (Ramb)	Nil	
		<i>Brachythemis contaminate</i> (F.)	Ditch Jewel	
		<i>Crocothemis servilia</i> (Durry)	Rudy Marsh Skimmer	
		<i>Crocothemis erythrea</i> (Brulle)	Broad Scarlet	
		<i>Orthetrum pruinosum</i> (Ramb.)	Crimson Tailed Marsh Hawk	
		<i>Orthetrum sabina</i> (Durry)	Green Marsh Hawk	
		<i>Rhyothemis variegata</i> (Linn.)	Common Picture Wing	
		<i>Tramea basilaris</i> Kirby	Red Marsh Trotter	
		<i>Tramea limbata</i> (Ramb.)	Black Marsh Trotter	
		<i>Tramea aurora</i> (Burm.)	Crimson Marsh Glider	
		Aeshnidae	<i>Anax parthenope</i> (Selys)	Yellow Ring Emperor
		Gomphidae	<i>Ictinogomphus rapax</i> (Ramb.)	Common Club tail
Zygoptera	Coenagrionidae	<i>Ceriagrion coromandelianum</i> (F.)	Coramandal Marsh Dart	
		<i>Ishnura senegalensis</i> (Ramb.)	Senegal Golden Dart	
		<i>Pseudagrion rubiceps</i> (Selys)	Saffron Faced Blue Dart	
		<i>Pseudogrion microcephalum</i> (Ramb.)	Blue Grass Dart let	
	Lestidae	<i>Lestes elatus</i> Hagen	Emerald Spread Wing	

The data indicates that the communities do not show a highly nested pattern ($T=36.07^0$ and $P= 3.86e-21$) and the matrix fill is only 46.3%. This is significantly a hot matrix. For matrix fills less than 50% a perfectly ordered matrix forms a concave meniscus in the upper left corner of the figure which is not observed in the present study. The Mean $T= 65.4^0$ for a Monte Carlo run count of 100 randomized matrix (Fig.1). Highly nested species were *Brachythemis contaminate* (♀), *Ishnura senegalensis*, *Orthetrum sabina*(♀),*Crocothemis*

possibility that in due course these idiosyncratic species may be lost forever from the regional biodiversity. These species can be identified by an idiosyncratic temperature higher than the systems temperature of the matrix (100⁰). However such species were not recorded and in some manner were disconnected from the local ecological system.

Odonates constitute an important segment of lakes ecosystems and their conservation is an important aspect of ecology. Identifying nested and idiosyncratic species can be done using the Nestedness pattern of assemblages. Monitoring lake ecosystems is one way of maintaining a balanced population of diverse Odonata species, and in turn a balanced ecosystem.

CONCLUSION

Nestedness pattern of Odonata insect communities around Hadhinaru Lake of Mysore was not quite significant. The matrix temperature was hot (36.7⁰). Secondly the matrix was not fully packed and idiosyncratic species (unexpected) were in higher numbers. Species that were nested were probably endemic. Allocathonous(disturbances due to human intervention) activities around the lake usually alter lake communities. The nestedness software serves as a basic tool to understand biodiversity of insects. Conserving insect diversity often provides a healthy aquatic ecosystem.




Fig.1. Nestedness assemblages for Odonata species

REFERENCES

1. Atmar. W., and Patterson.B.D.(1993). The measure of order and disorder in the distribution of species in fragmented habitats. *Oecologia*.96:373-382.
2. Atmar.W. and Patterson. B.D.(1995). The nestedness temperaturecalculator: a visual basic program, including 294 presence-absence matrices.AICS Res. University Park, New Mexico and the Field Mus. Chicago
3. Brown.J.H.(1995)Macro ecology. Chicago IL. University of Chicago Press.
4. Cook.R.R., and Quin.J.F. (1995).The influence of colonization in nested species sub-sets *Oecologia*: Vol.102. 413-424
5. Conard .K.F.,Wilson.K.H.,Harvey.I.F.,Thomas.C.J., and Sherratt.T.N(1989). Dispersal characteristics of seven Odonate species in an agricultural land scape.*Ecogeography*.Vol.22.524-531.
6. Emiliyamma.K.G., Radhakrishnan.C. and Palot.M.J. (2005). Common dragonflies and damselflies of Kerala. *Zoological Survey of India*.
7. Foote.A.L., Christine.L., and Hornung.R.(2005). Odonates as biological indicators of grazing effects on Canadian Praire Wetlands. *Ecol. Entomology*. Vol.30.273-283.
8. Fraser.F.C.,(1933). The fauna of British India, including Ceylon and Burma. *Odonata* Vol.I.Taylor and Francis Ltd. London.



Anax parthenope (Selys)



*Lestes elatus*Hagen(♂)



Brachythemis contaminata (F.)(♂)



Lestes elatus Hagen (♀)



Brachythemis contaminata (F.)(♀)



Crocothemis erythrea(Brulle) (♂)



Crocothemis servillia (Drury)(♂)



Crocothemis erythra (♀)



Crocothemis servillia(Drury)(♀)



Tramea limbata(Ramb.)




Rhyothemis variegata(Linn.)



Trithemis aurora(Burm)

Plate 1



Tramea basilaris(Kirby)(♀)



Ictinogomphus rapax(Ramb.)



Orthetrum sabina(Drury)(♂)



Orthetrum prunosum(Ramb.)(♀)



Orthetrum prunosum(Ramb.)(♂)



Ischnura senegalensis(Ramb.)



Pseudogrio microcephalum(Ramb.)



Ceriagrion coromendelianum(F.)



Pseudagrion rubriceps(Selys)



Ischnura elegans



Acisoma panorpoides Ramb



Brachydiplax sorbina (Ramb.)

Plate 2

9. Fraser.F.C. (1934). The fauna of british India, including Ceylon and Burma.Odonata Vol.II.Taylor and Francies Ltd. London.
10. Heino.J and Muota.T.(2005).Highly nested snail and calm assemblages in Boreal lake littorals. Roles of isolation, area, and habit suitability. Ecoscience. Vol. 12.141-146.
11. Levins.S.A.(1992).The problem of pattern and scale in ecology. Ecology. Vol.73.1943-1967.
12. Mellanby.M(1963). Animal Life in Freshwater. 6th Edition. Chapman and hall Ltd. London,UK.
13. Moore. N.W.(1953). Population density in adult dragonflies(Odonata:Anisoptera). J. Animal ecology.Vol 22.344-359.
14. Patterson.B.D.(1991). The integral role of biogeographic theory in the conservation of tropical forest diversity(eds. Mares.M.A. and Schmidly.D.J latin American Mammalogy. History, Biodiversity and Conservation. University of Okalahama Press. Norman.124-149.
15. Pennak.R.W.(1978). Freshwater invertebrates of the U.S. 2nd Edition. John Wiley and Sons.New York, USA.
16. Soinien.J.,Heino.J.,Kokocinski.M, and Muokta.T.(2009). Local regional diversity relationship varies with spatial scale in lotic distoms.J.Biogeog. Vol.36.720-727.
17. Subramanian.K.A. (2005).Dragonflies and damselflies of Peninsular India-A field Guide. Gadgil.M.(ed.). Indian Institute of Science and Indian Academy of Sciences.Bangalore.

18. Tonapi.G.T.(1980).Freshwater Animals of India:An Ecological Approach.Oxford & IBH Publishing Co. New Delhi
19. Ward.H.B.and Whipple.G.C.(1959).Freshwater Biology. Edmondson.W.T.(ed.) 2nd Edition. John Wiley and Sons. New York, London.
20. Werner .E.E. and Peacor.S.D.(1990). A review of trait-mediated indirect interactions. Ecology Vol.34.1083-1100. (1990)
21. Wright. D.H., Patterson. B.D., Mikkelson .G.M., Cutler. A.H. and Atmar. W. (1998). Acomparative analysis of nested subset patterns of species cosmopolitan. Oecologia Vol.113.1-20.